


ATP II – Cosmic Rays (WS22/23)

Air Shower Detection



Recap: Air Showers

Nucleus-induced Shower

• number of muons: $N_\mu = A \cdot \left(\frac{E_0/A}{\varepsilon_\pi} \right)^\beta = \left(\frac{E_0}{\varepsilon_\pi} \right)^\beta A^{1-\beta}$ e.g. $M=50, \beta=0.9$
 $\rightarrow N_\mu(56)/N_\mu(1) = 56^{0.1} = 1.5$

• energy in γ, e^+, e^- : $E_{em} = E_0 - N_\mu \cdot \varepsilon_\pi$ e.g. $E_0 = 10^{20}$ eV, $M=50, A=1 \Rightarrow E_{em}/E_0 \approx 91\%$
 $A=56 \Rightarrow E_{em}/E_0 \approx 86\%$

• number of γ, e^+, e^- : $N_{em} = E_{em} / \varepsilon_{crit} = \frac{E_{em}}{E_0} \cdot N_{em}^0$

• Shower maximum from 1st interaction:

$$X_{max}^A = \lambda_p + X_0 \ln \left(\frac{E_0}{2 \cdot A \cdot M \cdot \varepsilon_c} \right)$$

$A \cdot \frac{1}{3} M \pi^0$ with $E = E_0/A/M \Rightarrow 2 \cdot A \cdot \frac{1}{3} M \gamma$ s with $E_\gamma = E_0/A/M/2$

$$\underline{X_{max}^A(E) = X_{max}^P(E/A)}$$

how to measure the CR mass + energy with air showers


measurement	primary CR	detector
N_e, N_μ	$\leftrightarrow E_0, A$	SD
X_{max}, E_{em}	$\leftrightarrow E_0, A$	FD/CD
E_{em}, N_μ	$\leftrightarrow E_0, A$	RD+SD
$N_e, N_\mu, X_{max}, E_{em}$	$\leftrightarrow E_0, A$	SD+FD

} "hybrid"

overconstrained $\Rightarrow$ check hadronic interaction models

(SD: surface detector (particles)
 FD: fluorescence detector
 CD: Cherenkov detector
 RD: radio detector)

Air Shower Detection



Air Shower Detection








Figure 2

Average (a) lateral and (b) longitudinal shower profiles for vertical, proton-induced showers at 10^{19} eV. The lateral distribution of the particles at ground is calculated for 870 g cm^{-2} , the depth of the Pierre Auger Observatory. The energy thresholds of the simulation were 0.25 MeV for γ and $e^\pm$ and 0.1 GeV for muons and hadrons.


E and A

$\theta = 0^\circ$, $X = 800 \text{ g/cm}^2$





* E_{em} in lecture 10

Particle Detectors




Telescope Array "TA"

500 SCWTs
1.2 km spacing
 $A = 700 \text{ km}^2$



Pierre Auger Observatory

1600 WCDs
1.5 km spacing $> 10^{18} \text{ eV}$
 $A = 3000 \text{ km}^2$



both at $X_{\text{ground}} \approx 850 \text{ g/cm}^2$

Telescope Array: scintillator as particle detector




Signal

$$S \sim N_{ch} = N_{\mu} + N_e$$

($\gamma \rightarrow e^+e^-$ conversion)

$$\text{Area} \sim \cos \theta$$




Pierre Auger Observatory: Water Cherenkov detectors (WCD)



area $\sim$ const (volume detector, $h \sim r$)

12t ultrapure water




Cherenkov threshold: $\sim 0.8 \text{ MeV } e^-$
 $\sim 160 \text{ GeV } \mu^-$

Signal $\frac{dN_c}{dE} \sim E \rightarrow$ more signal for muons that traverse full det.

$$S = a \cdot N_\mu + b \cdot N_{e/\gamma}$$

IceTop @ IceCube




"Ice-Cherenkov" detectors



WCDs

Shower Front / arrival times

→ arrival direction



Type-I Station

Labels in the diagram include:


- Detector Station
- HV- and Signal Connectors
- Photomultiplier Tube
- Light Guide
- Pressure Balance Vessel (Ar filling)
- e/gamma-Detectors Liquid Scintillator
- Lead Absorber (10 cm)
- Iron Absorber (4 cm)
- Concrete
- Muon-Detectors Plastic Scintillator

Dimension: 2400mm

$$\rightarrow N_{ch} = N_{e^{+}e^{-}} + N_{\mu}$$




→ 20 attenuation lengths shielding

→ N_μ $E_\mu > 250 \text{ MeV}$



T.Stanev, "High Energy Cosmic Rays" (2010)

N_e/N_μ @ Auger



"AugerPrime" → additional scintillator on each station (since 2022)

schematically:


$$S_{\text{SWR}} \sim N_\mu + N_e$$

$$S_{\text{WCO}} \sim a N_\mu + b N_e$$

$$\begin{pmatrix} S_{\text{SWR}} \\ S_{\text{WCO}} \end{pmatrix} \sim \begin{pmatrix} 1 & 1 \\ a & b \end{pmatrix} \begin{pmatrix} N_\mu \\ N_e \end{pmatrix}$$

⇒ N_e and N_μ



highest energy event measured with an PD



PDG 2022


NEW HIGHEST EVENT DETECTED BY TA

- * 2021/05/27 10:35:56.47, No FD observation
- * $E = 243.6 \pm 10.7 \text{ EeV}$, $\theta = 38.6^\circ$, $\phi = 206.8^\circ$ - **Preliminary**
- * ($E = 242.8 \text{ EeV}$ with the atmospheric energy correction) - **Preliminary**




23

DIRECTION IN THE SKY-MAP



J. Matthews, TA Coll. 2022

Fluorescence Telescopes



Air Fluorescence

- N_2 excitation by charged particles


- isotropic emission !!

- fluorescence yield $Y \sim f(S, T, H) \cdot \frac{dE}{dx}$
density S
temperature T
humidity H
energy deposit in atmosphere

- rule of thumb: $\approx 3-4$ photons/m/particle
 $\approx 30W$ light bulb

- precise measurement in lab

emission in UV ($\sim 10ns$)



show visible up to $30km @ 10^{10} eV \Rightarrow 3000 km^2$


$45 km @ 10^{20} eV \Rightarrow 6000 km^2$

but: duty cycle $\lesssim 15\%$ (moon, sun, clouds, thunderstorms...)

"Schmidt optics"




6 telescopes per building
4 buildings at edge of array (looking inward)





spherical mirror

camera




440 hexagonal PMTs



⇒ see separate file for animation

Longitudinal profile






Nucl.Instrum.Meth.A 798 (2015) 172

Gaisser-Hillas function (GH function):

$$N_{fluo} \sim \frac{dE}{dX} \sim N_e(X) = N_{max} \left(\frac{X - X_1}{X_{max} - X_1} \right)^{(X_{max} - X_1)/\Lambda} e^{-\frac{X - X_{max}}{\Lambda}}$$

$$(X_1 \approx 0 \text{ g/cm}^2, \Lambda \approx 60 \text{ g/cm}^2)$$

highest energy fluorescence detector event



Fly's Eye Coll., ApJ 441 (1995) 144


see also T.Fitoussi et al, JCAP 01 (2020) 042

$$RA = 85.2^\circ, DEC = 48^\circ$$

$$E = (320 \pm 90) \cdot 10^{20} \text{ eV} \quad (51 \text{ Joule!})$$

$$X_{\text{max}} = (815 \pm 60) \text{ g/cm}^2 \quad (\text{compatible with } p/\text{Fe}/8)$$


“Hybrid Detection” (e.g. fluorescence and particles)





“Hybrid Detection” (e.g. fluorescence and particles)


Angular event display



Colors: time
Size: $\log(\text{signal})$






S_{1000}








S_{1000}



Non-Imaging^{*} Cherenkov Detectors



^{*} See summer semester for Cherenkov telescopes





$$N_g \sim N_e \sim E_{em}$$

"Calorimetric measurement"

(X_{max} from lateral shape and time dispersion)

TUNKA army




85m
spacing

51° 48' 35" N
103° 04' 02" E
675 m a.s.l.



Radio Detectors





$\lambda > \text{shower front thickness}$

$\Rightarrow f \lesssim 100 \text{ THz}$

shower $\approx$ point charge $\Rightarrow$ radial $\vec{E}$ from Q

$\vec{E} \perp \vec{B}$ from $\vec{I}$



T. Huege, Phys.Rept. 620 (2016) 1

geomagnetic effect

$>$

in air!


(Askaryan dominates in solids)

charge excess / Askaryan effect

$\approx 10\%$ in air

$$\vec{E} \sim N_e \Rightarrow \text{radiated power} \sim N_e^2 \sim E^2$$

Auger Coll., PRL 116 (2016) 241101



2p1c



LOPES@KASCADE



SALLA@Tunka




LPDA@Auger




SALLA@Auger


Some Results (E and A)





Energy Spectrum




Energy Spectrum



Mass Composition




Mass Composition



elongation rate

$$D_{10} > D_{10}^{\text{had}}$$

$$D_{10} < D_{10}^{\text{had}}$$



Detector Score Card (UHE)

	EAS variable	detector density	duty cycle	cost/unit	model dependence	maintenance/calibration
particle	N_E/N_M	$\approx 1/\text{km}^2$	$\approx 100\%$	low... medium $N_E \sim N_M$	high	low
fluorescence	$E_{\text{em}}/X_{\text{max}}$	$\approx 1/2000 \text{ km}^2$	$\leq 15\%$	high	low	high
radio	$E_{\text{em}}/X_{\text{max}}$	$1 \dots \gtrsim 100/\text{km}^2$ E_{em} with X_{max}	$\approx 100\%$	low... medium electronics!	low	low
Cherenkov	$E_{\text{em}}/X_{\text{max}}$	$\gtrsim 100/\text{km}^2$	$\leq 15\%$	low... medium	low	medium... high