

MODERN PHYSICS

Instructor: Prof. Bernd Pilawa (bernd.pilawa@kit.edu)

Winter Semester 22/23

Exercise 5

§ Wave-Particle Dualism §

Problem 1: Ruby laser

The ruby laser is a solid state laser consisting of an Al_2O_3 rod doped with about 0.05% of Cr. On one end the rod has a highly reflecting mirror surface and a semi-transparent one on the other. A pulsed xenon flash bulb is used to excite the chromium atoms from a ground state $|1\rangle$ to an excited state $|3\rangle$ which then quickly decays into a metastable state $|2\rangle^*$. Neglecting the excited state lifetime, the three level system is characterized by the following rate equations:

$$\begin{aligned}\frac{dN_1}{dt} &= -W_p N_1 n_\gamma + \frac{N_2}{T_{21}} \\ \frac{dN_2}{dt} &= +W_p N_1 n_\gamma - \frac{N_2}{T_{21}} \\ N &= N_1 + N_2 + N_3 \approx N_1 + N_2,\end{aligned}\tag{1.1}$$

where W_p is the stimulated transition probability resulting from the pumping process, n_γ is the number of pump photons, N_i are the state populations and T_{ij} the corresponding state decay times.

- a) Show that the population difference $\Delta N = N_1 - N_2$ in the steady state is given by the following equation:

$$\Delta N = N \cdot \frac{\frac{1}{T_{21}} - W_p n_\gamma}{\frac{1}{T_{21}} + W_p n_\gamma}.\tag{1.2}$$

- b) Under what conditions can a population inversion ($N_1 - N_2 < 0$) be achieved?
c) Why is the three-level system inefficient?


Problem 2: Compton effect

The Compton effect refers to the decrease in energy of a photon which is scattered inelastically by a charged particle. A.H. Compton used photons with a wavelength of $\lambda = 0.0711$ nm for his experiments (Molybdenum K_α -line).

- a) How large is the energy of these photons?
b) Calculate the wavelength of the photons which are scattered at free electrons under an angle of $\theta = 180^\circ$ (\Rightarrow in the direction of the source).
c) How large is the energy of one of these reflected photons?

Problem 3: Electron diffraction I: De Broglie wavelength

In an evacuated high vacuum chamber electrons are emitted from a hot cathode and accelerated through an electric potential difference. The resulting electron beam propagates through a polycrystalline graphite thin film. Behind the graphite layer, a hemispherical fluorescent screen with a radius of $R = 6.5$ cm (fig. left) is excited via the incident electrons and luminous effects in the form of fluorescent fringes can be observed.



- Explain why the following pattern can be seen (fig. right).
- Calculate the *de Broglie* wavelength of the electrons which are accelerated by a voltage of $U = 10$ kV.
- Determine the angles at which the electrons are scattered into the rings of the first two diffraction orders. The diameters of the first two diffraction rings are $D_1 = 1.5$ cm and $D_2 = 2.6$ cm (fig. right).
- Calculate the distances of the lattice planes within the crystallite of the graphite layer which correspond to the diffraction pattern/rings.

Problem 4: Electron diffraction II: Double slit

The setup from problem 3 is now modified. The emitted electrons are accelerated through an electric potential difference of $U = 100$ V and transmitted through a sample with two vertical slits. The slits have a distance of $d = 1$ μm between each other and a width of $a_1 = 0.5$ μm .

- Calculate the *de Broglie* wavelength of the electrons.
- Sketch the expected intensity pattern which is observed at a flat fluorescent screen at a distance of $l = 1$ m behind the double slit. What is the distance between the diffraction orders assuming incident plane waves?
- What changes when the potential difference is halved? What is the result of using a sample with the same slit separation, but with a slit width of $a_2 = 0.25$ μm ?