

# MODERN PHYSICS

Instructor: Prof. Bernd Pilawa (bernd.pilawa@kit.edu)

Winter Semester 23/24

## Exercise 11

### § Atoms & Solids §

#### Problem 1: Internal magnetic fields

From the point of view of an electron with orbital angular momentum  $\vec{L}$  the positively charged nucleus revolves around the electron. The motion of the nucleus produces an internal magnetic field parallel to the electron's orbital angular momentum. The intrinsic spin magnetic moment  $\vec{\mu}_s$  of the electron interacts with this internal magnetic field. The potential energy  $E_B$  of a magnetic moment  $\vec{\mu}$  in a magnetic field  $\vec{B}$  is given by the following equation:

$$E_B = -\vec{\mu} \cdot \vec{B}. \quad (1.1)$$

Since  $\vec{\mu}_s$  is proportional to the spin of the electron  $\vec{S}$  and since  $\vec{B}$  and  $\vec{L}$  are proportional for a given orbit, we can deduce that the potential energy associated with spin-orbit coupling is of the form:

$$E_S \propto \vec{S} \cdot \vec{L} \quad (1.2)$$

- Estimate the strength of the internal magnetic field due to the electron's orbital motion which results in the two sodium D-lines with  $\lambda = 588.995 \text{ nm}$  and  $\lambda = 589.592 \text{ nm}$ .
- Estimate the strength of the internal magnetic field due to the electron's orbital motion which results in the  $\lambda = 766.41 \text{ nm}$ - and  $\lambda = 769.90 \text{ nm}$ -lines observed in the  $L = 1$  to  $L = 0$  transition in potassium.

#### Problem 2: The Landé factor

The Landé factor  $g_J$  is the proportionality constant that relates the observed total magnetic moment  $\mu$  of an electron, resulting from both spin and orbital angular momentum, to its total angular momentum  $\vec{J}$ :

$$\vec{\mu} = -g_J \mu_B \frac{\vec{J}}{\hbar}. \quad (2.1)$$

With the g-factor for the electron is taken to be approximately  $\approx 2$ , we can express the Landé factor via the following equation:

$$g_J = \frac{3}{2} + \frac{s(s+1) - l(l+1)}{2j(j+1)} \quad (2.2)$$

- Show in a vector diagram that  $\vec{\mu}$  and  $\vec{J}$  are not parallel.
- Derive the Landé factor given in eq: g-factor by calculating the projection of  $\vec{\mu}$  on the vector  $\vec{J}$ .

*Hint:* The required projection term is given by the following equation:

$$\frac{\vec{\mu} \cdot \vec{J}}{|\vec{J}|} = -\mu_B \frac{\vec{J} \cdot \vec{J} + \vec{J} \cdot \vec{S}}{\hbar \cdot |\vec{J}|} = -\mu_B \sqrt{j(j+1)} g_J. \quad (2.3)$$

### Problem 3: Hexagonal lattice

The primitive translation vectors of the hexagonal Bravais lattice are given as:

$$\vec{a}_1 = \frac{\sqrt{3}a}{2}\vec{x} + \frac{a}{2}\vec{y}, \quad \vec{a}_2 = -\frac{\sqrt{3}a}{2}\vec{x} + \frac{a}{2}\vec{y}, \quad \vec{a}_3 = c\vec{z}. \quad (3.1)$$

- a) Show that the volume of the primitive cell spanned by the above vectors is  $\frac{\sqrt{3}a^2c}{2}$ .
- b) Show that the primitive translation vectors of the reciprocal lattice are the following:

$$\vec{b}_1 = \frac{2\pi}{\sqrt{3}a}\vec{x} + \frac{2\pi}{a}\vec{y}, \quad \vec{b}_2 = -\frac{2\pi}{\sqrt{3}a}\vec{x} + \frac{2\pi}{a}\vec{y}, \quad \vec{b}_3 = \frac{2\pi}{c}\vec{z}. \quad (3.2)$$

- c) Describe and sketch the first Brillouin zone of the hexagonal lattice.

### Problem 4: Laue & Bragg Condition

The crystal structure of a lattice can be determined by scattering of particles (photons, neutrons, electrons, etc.)

- a) Write up the condition for constructive interference according to Bragg.
- b) Write up the condition for constructive interference according to Laue.
- c) Give the relation between the basis vectors of the Bravais lattice and the reciprocal lattice.
- d) What is the geometrical meaning of a vector of the reciprocal lattice?